

M. Urrutia Pereira¹; L. Pitrez Mocelin²; HJ. Chong-Neto³;
H. Badellino⁴; V. Riquelme Martinez²; P. Oliveira Lima²;
R. Coelho Figueiredo⁵; O. Calderon Llosa⁶; JI. Larco Sousa⁶;
M. Soria⁷; A. De Souza Machado⁸; R. De Carvalho Baldaçara⁹;
D. Moura¹⁰; MS. Repka De Ramirez¹¹; MI. Rojo¹²; G. Lopéz Pérez¹²;
V. Acosta¹³; M. Paz Serrate¹⁴; EA. Costa Jordão¹⁵;
M. Valentín Rostan¹⁶; M. Calvo Gil¹⁷; P. Latour Staffel¹⁸; D. Sole¹⁹
¹Federal University of Pampa (UNIPAMPA), Uruguaiiana, Brazil; ²Federal
University of Pampa, Uruguaiiana, Brazil; ³Federal University of Parana,
Curitiba, Brazil; ⁴Clinica Regional del Este, San Francisco, Argentina;
⁵Hospital Regional De Augustinópolis, Tocantins, Brazil; ⁶Peruvian
Society of Allergy Asthma and Immunology, Lima, Peru; ⁷H.I.G.A. SAN
MARTIN LA PLATA, La Plata, Argentina; ⁸Federal University of Bahia –
Faculty of Education, Salvador, Brazil; ⁹State University of Tocantins –
UNITINS, Palmas, Brazil; ¹⁰IPS-Geriatric Hospital Gerardo Buongermini,
Asunción, Paraguay; ¹¹National University of Asuncion, San Lorenzo,
Paraguay; ¹²Mexican Allergy Society, Mexico City, Mexico; ¹³Hospital
Pediátrico Juan Pablo II, CUF, Argentina; ¹⁴Bolivian Allergy Society,
La Paz, Bolivia; ¹⁵Brazilian Alergic Society, Rio de Janeiro, Brazil;
¹⁶Allergy Medical Institute British Hospital, Punta del Este, Uruguay;
¹⁷Universidad Austral de Chile, Valparaíso, Chile; ¹⁸Universidad
Nacional Pedro Henríquez Ureña, Santo Domingo, Dominican Republic;
¹⁹Federal University of São Paulo, São Paulo, Brazil
*Presenting author: M. Urrutia Pereira