

I. Cherrez-Ojeda^{1,2}; A. Kasperska-Zajac³; R. Meshkova⁴; D. Fomina⁵; D. Unal⁶; S. Demir⁶; N. Teovska Mitrevska⁷; Ü. Şahiner; D. İlgün Gürel⁸; E. Özkaya⁶; O. Calderon Llosa⁹; L. Escalante¹⁰; A. Maldonado¹¹; E. Escalante¹¹; M. A. Touriz¹⁰; M. Faytong-Haro^{1,2}; M. F. Osorio^{1,2}; P. Staubach¹²; L. Herzog¹³; T. Roth¹⁴; K. Kulthanan¹⁵; C. Mann¹⁶; D. Cevallos-Levicek^{1,2}; K. Robles^{1,2}

¹Universidad de Especialidades Espíritu Santo, Samborondón, Ecuador;

²Respiralab Research Group, Guayaquil, Ecuador; ³University of Silesia

in Katowice, Katowice, Poland; ⁴Smolensk State Medical University,

Smolensk, Russia; ⁵city clinical hospital 52, Moscow, Russia; ⁶Istanbul

University Faculty of Medicine, İstanbul, Türkiye; ⁷Remedika General

Hospital, skopje, Greece; ⁸Hacettepe university, Ankara, Türkiye;

⁹Clinica SANNA El Golf, Lima, Peru; ¹⁰University of Guayaquil (UG),

Guayaquil, Ecuador; ¹¹Ephora Research Group, Guayaquil, Ecuador;

¹²Charité-Universitätsmedizin Berlin, Berlin, Germany; ¹³Fraunhofer

Institute for Translational Medicine and Pharmacology ITMP, Berlin,

Germany; ¹⁴Henry Ford Hospital, Miami, USA; ¹⁵Mahidol University,

Bangkok, Thailand; ¹⁶University Medical Center Mainz, Mainz,

Germany

*Presenting author: I. Cherrez-Ojeda
